

Prescribing Pattern of Antibiotics in Pediatric Hospital of Kathmandu Valley

Palikhe N^a

Abstract

Introduction	Antibiotics are among the most commonly prescribed drugs in pediatrics. Because of an overall rise in health care costs, lack of uniformity in drug prescribing and the emergence of antibiotic resistance, monitoring and control of antibiotic use are of growing concern and strict antibiotic policies should be warranted. Before such policies can be implemented, detailed knowledge of antibiotic prescribing practice is important.
Objective	The main objective is to study the prescribing practice of antibiotic in hospital in-patient.
Methods	A retrospective study of one and half month's duration was undertaken during November- December of 2003. A total number of 121 patients were taken for the study.
Results	The average number of drugs per patient was 5.01±1.36 and antibiotics per patient was 2.41±1.02. Antibiotics were prescribed at least once for 93 percent of hospitalized children although only few of the patients receiving antibiotics had a proven bacterial infection. Among 121 patients clinically diagnosed with infectious diseases and treated with antibiotics, specimens were taken for culture in only 24 cases i.e. (19.8%) to identify pathogenic organisms. Only 13 specimens showed positive culture results. Infants less than 1 year received antibiotics more frequently than 1-5 and 5-12 years (40, 31 and 29%, P<0.001). Seventy-five percent of the total antibiotics were administered parenterally. Cephalosporin was the top most frequently prescribed antibiotics followed by penicillin group. Significant difference was found between age group of patient and disease encountered (P<0.000).
Conclusion	The fact that children below 1 year or infants are at special risk of receiving multiple courses of antibiotics, together with the knowledge that antibiotic resistance develops in this setting; suggest that strategies to control antibiotic use should focus on these patients' populations.
Key words	Antibiotic, Inpatient and Pediatrics.

Introduction

Paediatrics is the branch of medicines dealing with the development, diseases and disorders of children¹. Infancy and childhood is a period of rapid growth and development. Compared to adult medicine, drug use in pediatrics is not extensively researched and the range of licensed drugs in appropriate dosage form is limited. Drug therapy is considered to be major component of paediatric management in health care setting like hospital. Effective medical treatment of a pediatric patient is based upon an accurate diagnosis and optimum course of therapy, which usually involves a medication regimen.

Infants and children are among the most vulnerable population groups to contract illnesses. The use of antimicrobial agents, especially antibiotics has become a routine practice for the treatment of paediatric illnesses^{2,3}. The key role of antibiotics for the treatment of infectious diseases that are prevalent everywhere in developing countries may not be denied. However, there are also reports of an irrational use of

antibiotics^{4,5} which may even lead to infections that are worse than the originally diagnosed ones. The pediatricians and other medical personnel who provide health care for infants and children in developing countries confront a number of challenges during the day to day practice of medicine due to the shortage of appropriate drugs and other facilities.

The rising incidence of bacterial resistance to common antibiotics, particularly, multi- drug resistant pneumococci, has prompted the need to use antibiotics judiciously in paediatric practice. Many of the antibiotics are unnecessarily prescribed for viral infections such as common cold. In a Kentucky study, 60 percent of patients were prescribed antibiotics for the common cold⁶. In a Canadian study from Saskatchewan, 85% of antibiotics prescribed for respiratory tract infection in children less than 5 years of age were considered inappropriate.

The study on prescribing practice of antibiotic has already

Corresponding Address: Nami Palikhe, **PO Box:** 21266, Sinamangal (Duwakot), E-mail p_namu@hotmail.com, Department of Pharmacology, Kathmandu Medical College, Nepal

been carried out in different population in Nepal. Study conducted by Shankar⁷ et al in teaching hospital of western Nepal shows that 203 of the 687 patients admitted to the Internal Medicine ward were prescribed antibiotics during the study period. Ninety-eight patients were prescribed a single antibiotic, while 75 patients were prescribed two antibiotics. Twenty-three patients were prescribed 3 antibiotics, while 4 and 5 antibiotics were prescribed to 5 and 2 patients respectively. The most common condition for which an antibiotic was prescribed was chronic obstructive pulmonary disease (COPD) (22.7%). The other common conditions were lower respiratory tract infections (LRTI) (15.8%), urinary tract infections (UTI) (13.3%) and pneumonia (8.9%). Ampicillin, amoxicillin, metronidazole, ciprofloxacin and crystalline penicillin were the 5 most commonly prescribed antibiotics. Similarly study conducted by Rehana⁸ et al in B.P. Koirala Institute of Health Sciences, Dharan, Nepal shows that the average number of drugs per prescription was 5.26. Eighty-four percent of all prescriptions contained anti microbial agent (AMA). It was the most commonly prescribed (42.8%) group of drugs followed by anti-inflammatory and analgesics (13.1%). Cloxacillin was the most commonly prescribed AMA followed by ampicillin. Ciprofloxacin plus metronidazole combination was the most frequently used.

Study done by Kafle⁹ et al shows that 43 percent of cases was prescribed antibiotics. Similarly another study done in 1994 by Kafle¹⁰ et al shows that 55 percent of cases were prescribed antibiotics.

Appropriate drug utilization studies have been found to be crucial to evaluate whether drugs are properly used and utilized in terms of medical, social and economic aspects.

Several professional societies have issued guidelines designed to reduce the use of antibiotics world- wide by means of various control strategies¹¹⁻¹². Detailed knowledge of antibiotic prescription pattern is important before the policies and measures can be implemented.

Methodology

The prescribing pattern of antibiotic usage was investigated in general ward of major children hospital in Kathmandu valley. A prospective follow up study of one and half month's duration was undertaken during November- December of 2003. Medical records of paediatric in-patients of age falls between 1 month and 12 years were retrospectively reviewed. Neonates and patients from other ward are excluded. Every odd number of patients was selected for the study. Total sample size was 121. Patient characteristics such as age, sex, body weights, drugs taken before admission, duration of hospitalization were noted. Similarly admission and discharge diagnosis, and the condition of the patients on admission and discharge were recorded. Moreover, drug data, including name of the drug, dosage regimen (form, route, frequency and duration), and the date on which the pharmacotherapy was instituted, were recorded. Intravenous fluids, blood transfusion and nutritional preparations were not included in the study. Drug data and patient characteristic data were computed using Ms Excel and SPSS statistical package. The results were expressed as proportions or as means \pm standard deviation

(SD). Means have been compared by the Student's t-test. Paired sample t- test and independent sample t- test were used where appropriate. A difference was considered as significant if a P value was less than 0.05. The retrieved medical records contained most of the required information .

Result

The mean age of pediatric in- patient was 4.45 ± 3.9 . The male patients were 61.9 percent where as the female patients were 38.1 percent. Thirty-nine percentages of patients were exposed to herbal or modern drugs prior to admission in hospital. Previous drug history (PDH) couldn't be found out for 19 percent of patients. Initial diagnosis per patient was 1.07 where as final diagnosis per patient was 1.23. Initial diagnosis was different to the final diagnosis in 35 percent of patient

Table 1 : Pediatric in-patient characteristics

Parameters	%	(No)
Age		
1 month - 1 yr	28.9	(35)
1-5 yr	36.4	(44)
5-12 yr	34.7	(42)
Age (mean \pm SD years)	4.45 \pm 3.9	
Sex		
Male	61.9	(75)
Female	38.1	(46)
Cases with Previous Drug History		
Yes	38.8	(47)
No	42.1	(51)
Not Significant	19.0	(23)
Diagnosis / patient		
Initial	1.07	
Final	1.23	
No difference in the initial and final diagnosis	65.0	
Length of stay/patient (days)	7.29 \pm 4.26	

Table 2: Pediatric In-Patient's exposure to drug(s) and antibiotic(s)

Drug(s)/patient	%	(No)	Antibiotic(s) / patient	%	(No)
1	2	(3)	1	21	(25)
2	5	(6)	2	37	(45)
3	16	(19)	3	28	(34)
4	20	(24)	4	10	(13)
5	22	(27)	5	2	(2)
6	14	(17)	6	1	(1)
7	9	(11)	7	1	(1)
8	8	(10)			
>9	4	(4)			
Range	1-15		Range	1-7	
Mean \pm S.D	5.01 \pm 1.365		Mean \pm S.D	2.41 \pm 1.02	

The average number of drugs per patient was 5.01 \pm 1.36. More than 98 percent of the patients were exposed to, at least, two drugs. As high as 93 percent of patients were prescribed, at least, one antibiotic. The average number of antibiotic per patient was 2.41 \pm 1.02. The multiple antibiotics were 79 percent. Only one antibiotic was prescribed in 21 percent, two antibiotics in 37 percent (highest), three antibiotics in 28 percent, four antibiotics in 10 percent of and finally five or even more than five antibiotics were prescribed in about 4 percent of all cases. No single antibiotic was administered to the pneumonia (Pneu), urinary tract infection (UTI), pleural effusion and not diagnosed pediatric patient. All the antibiotics were in combination. Similarly multiple antibiotics treated 80

percent of cases of meningitis. Enteric fever (E. fever) was the diagnosis where 75 percent of cases were treated by single antibiotics.

Among all patient prescribed with antibiotics, pneumonia was the most common diagnosis. Pneumonia was the leading cause of hospital admission among infant. However, 5 percent of patients in hospital were not diagnosed during hospital stay. Enteric fever and other diseases were mostly found in age above 1 year. There was a significant difference between the age group of patient and disease encountered ($\chi^2=42.95$, $P=0.000$). (Table 3)

Table 3: Diagnosis pattern % (No) in different age group.

Diagnosis	(1 month-1 yr) Infant	1- 5 yr	5-12 yr	Total
Pneumonia	15.00 (18)	7.00 (8)	2.00 (3)	24.00 (29)
Meningitis	3.30 (4)	1.65 (2)	4.13 (5)	9.09 (11)
Enteric fever	0	5.00 (6)	2.00 (2)	7.00 (8)
Pyrexia of unknown origin	0	2.47 (3)	1.65 (2)	4.13 (5)
Not diagnosed	1.65 (2)	1.65 (2)	1.65 (2)	4.95 (6)
Combination	6.61 (8)	4.13 (5)	6.61 (8)	17.35 (21)
Pleural effusion	0	0.82 (1)	3.30 (4)	4.13 (5)
Others	2.47 (3)	12.39 (15)	13.22 (16)	28.09 (34)
UTI	0	1.65 (2)	0	1.65 (2)

Rabies encephalomyelitis, rheumatic chorea, occult sepsis, acute glomerulonephritis (AGN), rheumatic fever etc.

Combination: Acute gastro enteritis (AGE) with hypokalemia, AGE with Pneu, AGN, and hypertension encephalopathy, AGN+ UTI, E.fever with malaria with ulcer, Jaundice with Pneu, Pneu with hypertension etc.

Benzyl penicillin (C.P) was prescribed in highest percentage in pneumonia followed by cefotaxim and other antibiotics. Third

generation ceftriaxone was prescribed in highest percentage in case of meningitis followed by cefotaxim and other antibiotic

Table 4: Frequency (%) of individual antibiotic for specific diagnosis


Diagnosis	C.P	Cefo	Genta	Ceftria	Cefalex	Amox	Cefa	Oflo	A+C	Clox	Others
Pneumonia	26	19	13	2	12	10	7	0	6	2	3
Meningitis	14	17	10	21	0	3	0	0	0	4	31
Enteric fever	0	10	10	70	0	0	0	0	0	10	0
PUO	0	38	12	12	0	0	0	25	13	0	0
Pleural effusion	0	23	0	0	0	0	0	8	23	15	30
UTI	0	20	20	20	0	0	0	20	0	0	20
Not diagnosed	17	0	24	12	6	0	0	6	0	0	35
Combination	22	14	16	3	9	9	2	0	8	7	20
Others*	22	13	13	6	5	6	0	0	4	1	28

Others: Rabies encephalomyelitis, rheumatic chorea, occult sepsis, acute glomerulonephritis (AGN), rheumatic fever etc.

Combination: Acute gastro enteritis (AGE) with hypokalemia, AGE with Pneu, AGN, and hypertension encephalopathy, AGN+ UTI, E.fever with malaria with ulcer, Jaundice with Pneu, Pneu with hypertension etc.


Cefo: Cefotaxim; Genta: Gentamycin; Ceftria: Ceftriaxone; Cefalex: Cefalexin; Amox: Amoxicillin; Cefa: Cefaclor; Oflo: Ofloxacin; A+C: Ampicillin+Cloxacillin; Clox: Cloxacillin.

Figure 1:



Antibiotic of cephalosporin was the top most frequently prescribed antibiotic followed by penicillin group. Fluoroquinolones were prescribed by 6 percent.

Figure 2:.



In this study, 75 percent of the total antibiotics were administered parenterally. Drugs were prescribed in generic by 41 percent. Significant difference was found between the generic prescribing and brand prescribing of drug to children (P=0.000). Forty-seven percentage of the patients were prescribed at least one analgesic/antipyretic drug (AN/AP).

(n=121) for patient receiving at least one analgesic and antipyretic
(n₂ = 607) for others.

Among 121 patients clinically diagnosed with infectious diseases and treated with antibiotics, specimens were taken for culture in only 24 cases i.e. (19.8%) to identify pathogenic organisms. Only 13 specimens showed positive culture results. Sensitivity analysis was done in 6 cases (one E. fever and five UTI). In all cases (4-7) antibiotics were resistant. Null hypothesis set i.e. H₀ = 10 percent; 10 percent of patients are requested for culture of blood. Result from the analysis showed that 14.87 percent of patients were requested for blood

culture. The hypothesis is acceptable since no significant difference was found (P=0.0768) (Table 5).

Table 5: Antibiotic prescription and identification of pathogenic organism

Culture Information	Blood Culture	Urine culture	Both
Total culture-24	18	2	4
Culture +ve-13	7	2	1 BP +3 UP

BP: Both Positive
UP: Urine Positive

In this study out of 121, 11 of them had a record of adverse drug reaction associated with fever and rashes. Infant less than 1 year received antibiotics more frequently than 1-5 and 5-12 years. (40, 31 and 29 respectively, (P<0.001, P=0.000).

Sixteen percent of patients were in fair condition though their mean length of hospital stay was slightly greater than improved patient. Eight percent of patients were referred in different hospital according to the diagnosis (Table 6).

Table 6: Summary of treatment/Patient follow up evaluation

Patient condition	%	Mean hospital stay (days)
Improved	70	6.97
Fair	16	8.63
Discharge under request	3	7
Referred	8	8.6
Death	3	4.33

Discussion and Conclusion

The number of male patients was comparatively more than the number of female patients. Previous drug history was found in 38.8% of patients. This may be slightly above as there were still 19.0% of cases for which exact previous drug history couldn't be figured out properly. The average number of admission diagnosis per patient was 1.07. The average number of final diagnosis (1.23) was slightly greater than former ones. This may be due to proper diagnosis and the result of different investigations. Child may develop other complications due to various reasons such as nosocomial infection, complication of disease itself, complication due to inappropriate drug etc. The overall length of stay was less (7.29 days) compared to study done in Gondar hospital (24.9 days) and more than study done in Bahirdar hospital (4.8 days) of Ethiopia in 1993/1994 by Abula et al¹³. This may be due to available of better treatment facilities and technique than Gondar hospital. A high percentage of patients i.e. 93% was prescribed at least one antibiotic which was different than study done by Marlies et al¹⁴ i.e. 36 percent and Jason Hall¹⁵ i.e. 60.6 percent. Similarly study done by Bosu et al¹⁶ again showed the variation in average percentage of patients receiving at least one antibiotic, which was 41 percent, 45 percent, 79 percent and 98 percent in different health centers. It is not possible to draw any firm conclusion since the patients are not matched socio- economically. The morbidity pattern also may not be similar. Six and seven antibiotics were found to be prescribed in 1 percent of patient. This is due to requirement of more time to diagnose the patient and select the appropriate treatment. The prescription of antibiotics was based on mainly clinical judgement without investigation.

Benzyl penicillin and gentamycin / cefotaxim were found to be mostly used combination antibiotics in case of pneumonia. Similarly ceftriaxone in combination with chloramphenicol was used in case of meningitis. Enteric fever was generally treated by ceftriaxone. It is the only disease where exposure to combination antibiotic was found to be minimum.

Pneumonia was the most prevalent diagnosis among infant, which agrees with studies. Only 19.8 percent cases were demanded for the culture of blood or urine. Only 13 cases showed the positive results out of 24 so, only in a few patients therapy could be modified after culture results were available. Similar study was done by S. Hu et al¹⁷ where specimens were taken for culture in 8.4 percent. Specimens were not obtained in most cases in this hospital or patients were not asked for the culture report. It is very necessary to monitor appropriated clinical specimens have been obtained,

examined and cultured before antibiotic is initiated. The use of antibiotics is related to antibiotic resistance, side effects of drugs and health care costs. So, measures should be taken care to avoid the inappropriate use of antibiotics. Physicians must have a clear understanding of therapeutic use of antibiotics; they must be aware of the prevalence of various pathogens and resistance patterns in their hospital and exercise good judgment in selection empirical antibiotic regimens¹⁷. Sensitivity analysis was done in one case of enteric fever and five cases of urinary tract infection. In two cases, third generation of cephalosporin and expensive antibiotic "amikacin" were resistant. Ofloxacin, gentamycin, norfloxacin, nalidixic acid were resistant in all cases.

Infant less than 1 year received antibiotics more frequently was similar kind with that of the study done by Marlies et al¹⁴ which shows infant less than 2 years received antibiotics more frequently than older children (25 and 11% respectively, P=0.0256)

The excessive use of injectable is common in many developing countries¹⁸. In this study 75 percent of antibiotics were given by injection. It seems necessary for the paediatric patient to be treated by parenteral route of administration but consideration should be taken care for the syringes used to administer different antibiotic. Higher use of vitamins may be due to its use as placebo. Their benefits should also be viewed from their socio-economic and cultural aspect.

Over prescribing of analgesic/ antipyretic seems to be a problem. When the condition demands the use of such group of drugs, it is preferable to use paracetamol rather than aspirin and nimesulide as the former one is equally effective but has minimal adverse effects as compared to the later. It is noteworthy that nimesulide has been already banned for pediatrics. So, its use should be totally discontinued in pediatrics. The Register of adverse reactions of the Finnish National Agency for medicines (NAM) for the year 2000 shows that the majority of adverse drug reactions among non-steroidal anti-inflammatory diseases were associated with nimesulide; over half of the reports were associated with liver reactions. However nimesulide has never been marketed in some countries such as the USA and Australia. In South East Asia the drug enjoys variable regulatory status¹⁹.

The habit of recording adverse drug reaction must be encouraged at all level of health care institution. Fever was resulted due to ceftriaxone and rashes were found in case of patient treated by Ampicillin+Cloxacillin and ciprofloxacin. Adverse drug reaction was developed in ciprofloxacin prescribed patient due to the development of hypersensitivity reaction.

Cephalosporins were the top most used class of antibiotics in this study followed by penicillin. Among cephalosporins, third generation of ceftriaxone and cefotaxim were found to be mostly used. The use of fluoroquinolones by 6 percent of total antibiotics reminds that no quinolones were used by paediatric services because of their toxic effects in children below 14 years of age. Ciprofloxacin, as one of the frequently prescribed quinolone, deserves continued monitoring. Similar chloramphenicol was used mostly in case of meningitis with combination with cephalosporin. From the potentially serious adverse effects associated with the indiscriminate use of chloramphenicol, and a practice of empirical basis of prescription, the wide use of chloramphenicol may evoke some concern. In conclusion, the main challenges in prescription of antibiotics are to achieve a rational choice and appropriate use of antibiotics and to recognize their potential problems. Consequently, physicians must keep a clear understanding of need for microbiological diagnosis, use of antibiotics and make good judgment in clinical situations.

References

1. OP Ghai. Essential Pediatrics, fourth edition, published by *interprint A-16*, Naraina II, New Delhi-110 028, India.
2. Sanz EJ, Bergman U and Dahlstorm M. Paediatric drug prescribing. *Eur J clin pharmacol* 1989; 37:65-8.
3. Summers RS and Summers B. Drug prescribing in paediatrics. *Ann Trop paediatr* 1986; 6:129-33.
4. Principi N et al. Control of antibiotic therapy in paediatric patients. *Developmental pharmacology and therapeutics* 1981; 2:145-55.
5. Schollenberg E and Albritton WL. Antibiotic misuse in a paediatric teaching hospital. *Can Med Assoc J* 1980;122:49-52.
6. Dr. M. Ipp. Reduced antibiotic use in a Paediatric practice: Practical office strategies based on current evidence. Published in June 2000.
7. Shankar RV, et al. Prescribing patterns of antibiotics and sensitivity patterns of common microorganisms in the Internal Medicine ward of a teaching hospital in Western Nepal: a prospective study. *Ann Clin Microbiol Antimicrob* 2003; 2:7.
8. Rehana HS and Nagrani MA. A study on the drug prescribing pattern and use of antimicrobial agents at a tertiary care teaching hospital in eastern Nepal. *Indian Journal of Pharmacology* 1998; 30: 175-80.
9. Kafle KK. Pradhan YMS, Shrestha AD and Karki SB. Drug use in PHC facilities of Kathmandu. *Journal of the Institute of Medicine* 1992; 14:318-26.
10. Kafle KK. Prescribing and dispensing practices in PHC facilities of Terai districts of Nepal. *Journal of the Institute of Medicine* 1996; 18:61-6.
11. Kolar J, Kadakova E. Prescription of antimicrobial drugs to hospitalized children. *Ann Pharmacother* 1993; 26: 974-7.
12. Marr JJ, Moffet. Kunin CM. Guidelines for improving the use of antimicrobial agents in hospitals: a statement by the Infectious Disease Society of America. *J Infect Dis* 1988; 157; 869-76.
13. Abula T et al. Prescribing patterns of drugs in pediatric wards of three Ethiopian hospitals. Original 1993/1994.
14. Marlies A, Van H, Klarieke L. Antibiotic utilization for hospitalised paediatric patients. *International Journal of Antimicrobial Agents* 1998; 10:161-64.
15. Jason Hall BA. Paediatric prescribing in New Zealand. *NZFP* 2002; 29:14-8.
16. Bosu WK. Survey of antibiotic prescribing pattern in government health facilities of the Wasa-west district of Ghana. *East African Medical Journal* 1997; 74:139.
17. Suping Hu, Xiuheng Liu and Yan Peng. Assessment of Antibiotic Prescription in Hospitalized Patients at a Chinese University Hospital 2002.
18. Tomson G. Drug utilization studies in Sri Lanka- Towards an understanding of medicine in society. Thesis, Karolinska Institute, Stockholm, Sweden 1990.
19. Drug Bulletin of Nepal (DBN) 2003; 14: Royal pharmaceutical society 1996.