

Long-Term Effects of Topical Anti-Glaucoma Medications on Conjunctival Cell Morphology by Impression Cytology

Sarina Pachhai,¹ Madhu Thapa,² Pragati Gautam,² Nisha Sharma³

¹Department of Ophthalmology, Bheri Hospital, Nepalgunj, Nepal, ²B.P.Koirala lions centre for Ophthalmic Studies, Department of Ophthalmology, Maharajgunj Medical Campus, Institute of Medicine, Tribhuvan University, Nepal, ³Department of Pathology, Maharajgunj Medical Campus, Institute of Medicine, Tribhuvan University, Nepal.

ABSTRACT

Background: Glaucoma is a major cause of irreversible blindness worldwide. Reduction of intraocular-pressure (IOP) using topical anti-glaucoma medications remains the mainstay of treatment. However, prolonged use of these medications may induce adverse effects on the ocular surface.

Methodology: This cross-sectional observational study, conducted at the B.P. Koirala Lions Centre for Ophthalmic Studies, Maharajgunj Medical Campus from February 2023 to August 2024. Patients of primary open angle glaucoma receiving topical anti-glaucoma medications for at least three months were enrolled. Conjunctival- Impression Cytology samples were obtained and graded according to Nelson's grading system. Data analysis was done using SPSS version 16. A p-value <0.05 was considered statistically significant.

Results: 128 eyes from 128 glaucoma patients were analyzed, including 54 males (42.2%) and 74 females (57.8%), with mean age of 58.16 ± 14.44 years. Patients using prostaglandin-analogues demonstrated 39.3% Grade-2 and 27.3% Grade-3 changes. Duration of medication use exceeding three years was significantly associated with higher Nelson grades ($p < 0.05$). Older patients, particularly those aged 61–70 years, exhibited greater cytological changes ($p = 0.005$). A significant positive correlation was observed between duration of medication use and severity of conjunctival changes ($r = 0.597$, $p < 0.001$). Males exhibited more severe conjunctival changes than females ($p = 0.009$).

Conclusions: Prostaglandin-analogues showed most severe conjunctival cytological alterations. Longer duration of medication use, advancing age, and male gender were significantly associated with conjunctival changes.

Keywords: Anti glaucoma medications; conjunctival morphology; impression cytology, ocular surface disorder.

INTRODUCTION

Glaucoma is characterized by progressive damage to the optic nerve head, leading to irreversible blindness.¹ The number of individuals aged 40-80 years affected by glaucoma is expected to reach 111.8 million by 2040.^{2,3} Risk factors for glaucoma are family history, age, race and ethnicity, cardiovascular diseases and diabetes mellitus.^{4,6}

Use of topical anti glaucoma medications delays visual field loss by lowering intraocular pressure and reduces the risk of progression by 17% in patients.⁷The prolonged application of these medicine can lead to various ocular

surface disorder, manifesting as changes in conjunctival cell morphology.⁸ Ocular surface abnormalities will lead to dry eye related problems. That in turn will cause less compliance to anti-glaucoma drugs causing progression of glaucomatous optic neuropathy and ultimately results in glaucoma blindness.

Impression cytology is a non-invasive technique utilized to assess the morphology of conjunctival cells, goblet cell density and epithelial cell characteristics, which are indicative of ocular surface integrity.⁹ Nelson's grading system is a standardized method for categorizing these cytological changes.¹⁰

Correspondence: Madhu Thapa, B.P.Koirala lions centre for Ophthalmic Studies, Department of Ophthalmology, Maharajgunj Medical Campus, Institute of Medicine, Tribhuvan University, Nepal. Email: drmadhuthapa@gmail.com, Phone: +9779849537950.

This study aimed to investigate the long-term effects of various topical anti-glaucoma medications on conjunctival cell morphology using impression cytology.

METHODS

This was a hospital-based cross sectional observational study conducted at the B.P. Koirala Lions Centre for Ophthalmic Studies, Maharajgunj Medical Campus, Institute of Medicine, Kathmandu, Nepal. Ethical approval to conduct the research was taken from Institutional Review committee of Institution of Medicine, Tribhuvan University (Ref No: 426(6-11) E2 on 1/3/2023). Informed consent were taken from patients before enrolling in the study. All patients diagnosed with primary open-angle glaucoma (POAG) those receiving topical anti-glaucoma medications for a minimum duration of three months and who presented during from February 2023 to August 2024 were included in the study. A total of 128 eyes from 128 patients were enrolled. In cases where both eyes were treated, eyes which has been treated longer was included. Patients with obvious ocular surface disorders, history of ocular surface surgeries and cyclodestructive procedures in the past, long term use of any medications apart from anti glaucoma medications were excluded. More than 3 months uninterrupted use of antiglaucoma drugs is considered as long term use.

Patients using single drops were categorized according to the types of medications they are using. When multiple anti glaucoma medications are used in a single patient, it is considered as a combination therapy.

Conjunctival impression cytology was performed on the inferonasal bulbar conjunctiva of each enrolled eye, which was randomly selected. Cellulose acetate filter paper was used to collect epithelial cells from the ocular surface. The collected samples were then sent to pathology department Maharajgunj Medical Campus for cytological examination.

The severity of conjunctival cytological changes were assessed using Nelson's grading system.¹⁰ This system classifies conjunctival morphology based on characteristics such as goblet cell density, epithelial cell size and shape, and the presence of inflammatory cells. Grades typically range from zero (normal) to 3 described as below.

Grade 0: The epithelial cells are small and round with eosinophilic-staining cytoplasm. The nuclei are large, basophilic, with N/C ratio of 1/2. The goblet cells are

abundant, plump, and oval and have an intensely PAS-positive cytoplasm.

Grade 1: The epithelial cells are slightly larger and more polygonal and have eosinophilic-staining cytoplasm. The nuclei are smaller, with N/C ratio of 1/3. The goblet cells are decreased in number, however, they still maintain their plump oval shape, with an intensely PAS-positive cytoplasm.

Grade 2: The epithelial cells are larger and polygonal, occasionally multinucleated with variably staining cytoplasm. The N/C ratio is between 1/4 -1/5. The goblet cells are markedly decreased in number and are smaller, less intensely PAS-positive with poorly defined cellular borders.

Grade 3: The epithelial cells are large and polygonal with basophilic-staining cytoplasm. The nuclei are small, pyknotic, and completely absent from many cells. The N/C ratio is larger than 1/6. Goblet cells are completely absent.

Data were entered and analyzed in SPSS (version 16). Descriptive statistics were utilized to summarize the demographic characteristics of the study population, including age, gender, and laterality of the affected eyes. To investigate the association between different types of anti-glaucoma medications and the severity of conjunctival cytological changes, the Chi-square test was employed. Furthermore, Spearman's correlation coefficient was calculated to determine the relationship between the duration of medication use and the severity of conjunctival alterations, while the Mann-Whitney U test compared cytological changes between males and females. A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant for all analyses.

RESULTS

This study enrolled 128 eyes of 128 glaucoma patients. The study population included 74 (57.8%) female patients and 54 (42.2%) male patients. the mean age of the participants was 58.16 ± 14.44 years (ranging from 21 to 90 yrs).

Prostaglandin analogues were the most frequently prescribed anti glaucoma medication in this study, followed by combination therapy of various groups of drugs. Other medications included Beta-blockers, fixed formulation of alpha agonist and beta blocker, Alpha-agonists and Carbonic anhydrase inhibitors (CAI) as shown in Figure 1.




Figure 1. The Various Groups of Anti Glaucoma Medications Used.

Conjunctival cytological changes were assessed using Nelson’s grading system. The severity of these changes varied significantly across different types of medication. Patients using Prostaglandin analogues exhibited the highest severity of cytological changes, with 39.3% classified as Grade 2 and 27.3% as Grade 3. In contrast, Carbonic anhydrase inhibitors showed a lower incidence of severe changes, with 50% in Grade 0 and 41% in Grade 1 as shown in Table 1.

Table 1. Conjunctival Cytology Changes According to Nelson’s Grading Among Various Anti-glaucoma Drugs.

Medication Type	Grade 0 (%)	Grade 1 (%)	Grade 2 (%)	Grade 3 (%)	Total (n)	p-value
Alpha agonist	7.1	35.0	28.5	28.5	14	0.004
Alpha agonist + beta blocker fixed combination	20.0	40.0	40.0	0.0	15	
Prostaglandin analogue	12.0	21.0	39.3	27.3	33	
Carbonic anhydrase inhibitors	50.0	41.0	9.0	0.0	12	
Combination therapy	36.0	36.0	10.0	16.0	30	
Total	25.8	35.9	22.7	15.6	128	

(Chi Square test)

A statistically significant association was found between the duration of drug use and the severity of Nelson grades ($p < 0.05$). Specifically, patients who had been using anti-glaucoma medications for more than three years demonstrated higher Nelson’s grades, indicating more pronounced conjunctival alterations.

Table 2. The Cytological Grading in Relation to Duration of Use of Medications.

Duration of treatment (yrs)	Conjunctival cytological changes according to Nelson’s criteria				p-value
	Grade 1	Grade 2	Grade 3	Grade 4	
<1	2	1	0	0	0.9
1-3	13	9	0	1	0.1
>3	18	36	29	19	<0.05
Total	33	46	29	20	

(Chi-square test)

Spearman's correlation analysis revealed a significant positive correlation between the duration of medication use and the severity of conjunctival changes ($r = 0.597$, $p < 0.001$). This moderate to strong correlation suggests the cumulative impact of long-term anti-glaucoma therapy on ocular surface health.

Furthermore, older age groups, particularly those aged 61-70 years, (Chi-Square test, p value 0.005) were associated with greater cytological changes, suggesting that age is a contributing factor to ocular surface damage in glaucoma patients.




Figure 2. Bar Diagram Showing Conjunctival Morphological Changes Across Age Groups.

Males showed higher grades of conjunctival changes than females. The difference in Nelson's grading among males and females is statistically significant (Mann-Whitney U test, $p=0.009$)




Figure 3. The Distribution of Conjunctival Cytological Changes (Nelsons grading) Among Males and Females.

DISCUSSION

Primary treatment of glaucoma is using anti glaucoma medicine to control the IOP. This study provides valuable insights into the long-term effects of topical anti-glaucoma medications on conjunctival cell morphology in patients with primary open-angle glaucoma. Our findings highlights the significant impact of medication type, duration of treatment, patient age and gender on ocular surface health, as assessed by impression cytology using Nelson's grading system¹⁰.

The predominant use of prostaglandin analogues (PGAs) as the most frequently used medication is seen in current clinical practices, where PGAs are often considered first-line therapy due to their potent IOP-lowering efficacy and once-daily dosing.¹¹ Long term studies with latanoprost have not revealed any cardiac or pulmonary effects unlike timolol which may cause systemic symptoms like bradycardia, arrhythmia, congestive heart failure and syncope. So Better formulation, easy dosage, better compliance and less systemic side effects may explain the more use of prostaglandin analogue in glaucoma patients. However, our study also revealed that PGAs were associated with the highest severity of conjunctival cytological changes, with a substantial proportion of patients exhibiting Grade 2 and Grade 3 alterations similar to some published reports.¹² This finding is consistent with previous researchs indicating that PGAs, particularly those containing preservatives like benzalkonium chloride (BAK), can induce significant ocular surface toxicity, leading to inflammation, goblet cell loss, and squamous metaplasia.^{13,14} The chronic exposure to these agents can disrupt the delicate balance of the ocular surface, contributing to symptoms of ocular surface disease (OSD) and potentially affecting patient adherence to treatment.^{15,16} A report from Dogan et al has found that there is increased apoptosis of conjunctival epithelial cell examined by conjunctival biopsy in patients of glaucoma treated with anti-glaucoma medications.¹⁷ Similarly, Thakur et al have also observed the similar finding in eastern Nepal, where they had followed the newly diagnosed cases of glaucoma and ocular surface changes were seen within six months of treatment. Fixed combination therapy and prostaglandins drops showed a significant changes.¹⁸ A report from south India has shown similar that ocular surface disorder was prominent in patient on topical anti glaucoma drops and it was pronounced as the number drops and duration of treatment increased.¹⁹ Comparing the conjunctival morphology in people on anti glaucoma medications and those without these medicine significant changes were see as reported by

Pai et al.²⁰

The significant association between the duration of drug use and the severity of Nelson grades is a critical finding. Patients using anti-glaucoma medications for more than three years demonstrated more pronounced conjunctival alterations. Nelson' et al had also found the similar fact.²¹ This suggests a cumulative toxic effect of these medications on the ocular surface over time. This long-term impact highlights the importance of regular ocular surface assessment in glaucoma patients, especially those on prolonged therapy, to mitigate potential complications and improve their quality of life.²²

Furthermore, our study identified older age groups (61-70 years) as being more susceptible to greater cytological changes. This could be attributed to age-related physiological changes in the ocular surface, such as decreased tear production, reduced goblet cell density, and compromised epithelial barrier function, which may exacerbate the adverse effects of anti-glaucoma medications.²³ The combined effect of aging and medication induced toxicity contributes to the increased severity of OSD in elderly glaucoma patients.

When comparing the changes among males and females using anti glaucoma medication, the changes were more severe among males than females, which was unpredictable. Dry eye symptoms are reported more prevalent in females than males.²⁴ Studies in past have shown that females have more dry eye symptoms especially after the age of 40-50 years²⁵ due to hormonal changes, which is contradictory to our findings. This discrepancy in our study, might be explained by occupational or environmental exposures might be more in males.

Our results are largely in agreement with other studies that have utilized impression cytology to evaluate ocular surface changes in glaucoma patients. For instance, a study by Hong et al. similarly found a statistically significant difference in cytological changes among different anti-glaucoma drug groups.²⁶ While some studies, such as that by Turacli et al., reported no significant differences.²⁷ variations in study populations, methodologies, and duration of follow-up could account for these discrepancies. The reliability and non-invasiveness of impression cytology make it a valuable tool for monitoring ocular surface health in this patient population.

This study consisted of a small simple size and it is

a single center study so generalizability to wider population may not accurate. Secondly, age related ocular surface changes may be a confounding factor in older population of this study. We did not look into age changes in particular. Thirdly, preservative free anti-glaucoma medications are not available in Nepal. Otherwise, comparison with preservative free drops would have given a more conclusive result.

CONCLUSIONS

Long-term use of topical anti-glaucoma medications significantly affects conjunctival cell morphology, with the severity of changes influenced by type of medication, duration of treatment and patient age. Prostaglandin analogues were associated with the most severe ocular surface alterations, particularly in older patients and those on medication for more than three years. These findings emphasize the need for careful consideration of ocular surface health in the long-term management of glaucoma, advocating for regular monitoring and potentially the use of preservative-free formulations or alternative treatment strategies to minimize ocular surface toxicity.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Quigley HA. Glaucoma. *The Lancet* [Internet]. 2011 Apr [cited 2026 Feb 12];377(9774):1367-77. [Article]
2. Bourne RRA, Stevens GA, White RA, Smith JL, Flaxman SR, Price H, et al. Causes of vision loss worldwide, 1990-2010: a systematic analysis. *Lancet Glob Health*. 2013 Dec;1(6):e339-349. [Article]
3. Tham YC, Li X, Wong TY, Quigley HA, Aung T, Cheng CY. Global Prevalence of Glaucoma and Projections of Glaucoma Burden through 2040: A Systematic Review and Meta-Analysis. *Ophthalmology* [Internet]. 2014 Nov 1 [cited 2025 Jun 16];121(11):2081-90. [Article]
4. Kang JH, Wang M, Frueh L, Rosner B, Wiggs JL, Elze T, et al. Race/ethnicity in relation to incident primary open-angle glaucoma characterized by autonomously determined visual field loss patterns [Internet]. *medRxiv*; 2021 [cited 2026 Jan 13]. p. 2021.10.25.21265498. Available from: <https://www.medrxiv.org/content/10.1101/2021.10.25.21265498v1>
5. Grzybowski A, Och M, Kanclerz P, Leffler C, De Moraes CG. Primary Open Angle Glaucoma and Vascular Risk Factors: A Review of Population Based Studies from 1990 to 2019. *J Clin Med* [Internet]. 2020 Mar 11 [cited 2026 Jan 13];9(3):761. [PMC7141380]
6. Nislawati R, Taufik Fadillah Zainal A, Ismail A, Waspodo N, Kasim F, Gunawan AMAK. Role of hypertension as a risk factor for open-angle glaucoma: a systematic review and meta-analysis. *BMJ Open Ophthalmol* [Internet]. 2021 Sep 28 [cited 2026 Jan 13];6(1):e000798. [PMC8479953]
7. Schuster AK, Erb C, Hoffmann EM, Dietlein T, Pfeiffer N. The Diagnosis and Treatment of Glaucoma. *Dtsch Arzteblatt Int*. 2020 Mar 27;117(13):225-34.
8. Baudouin C, Labbé A, Liang H, Pauly A, Brignole-Baudouin F. Preservatives in eyedrops: The good, the bad and the ugly. *Prog Retin Eye Res* [Internet]. 2010 Jul 1 [cited 2026 Jan 9];29(4):312-34. [ScienceDirect]
9. Nelson JD. DRY EYE: DIAGNOSTIC TEST TEMPLATE.
10. Nelson JD. Impression Cytology. *Cornea* [Internet]. 1988 Mar [cited 2026 Feb 12];7(1):71. [Article]
11. Alm A, Stjernschantz J. Effects on intraocular pressure and side effects of 0.005% latanoprost applied once daily, evening or morning. A comparison with timolol. *Scandinavian Latanoprost Study Group. Ophthalmology*. 1995 Dec;102(12):1743-52. [Article]
12. Holló G. The side effects of the prostaglandin analogues. *Expert Opin Drug Saf*. 2007 Jan;6(1):45-52. [Article]
13. Jean MS, Brignole F, Bringuier A, Bauchet A, Feldmann G, Baudouin C. Effects of benzalkonium chloride on growth and survival of Chang conjunctival cells. *Invest Ophthalmol Vis Sci* [Internet]. 1999 Mar 1 [cited 2026 Jan 9]; [Article]
14. Sherwood MB, Grierson I, Millar L, Hitchings RA. Long-term morphologic effects of antiglaucoma drugs on the conjunctiva and Tenon's capsule

- in glaucomatous patients. *Ophthalmology*. 1989 Mar;96(3):327-35. [Article]
15. Walsh K, Jones L. The use of preservatives in dry eye drops. *Clin Ophthalmol Auckl NZ*. 2019;13:1409-25. [Article]
16. Asiedu K, Abu SL. The impact of topical intraocular pressure lowering medications on the ocular surface of glaucoma patients: A review. *J Curr Ophthalmol*. 2019 Mar;31(1):8-15. [Article]
17. Doğan AS, Orhan M, Söylemezoğlu F, Irkeç M, Bozkurt B. Effects of topical antiglaucoma drugs on apoptosis rates of conjunctival epithelial cells in glaucoma patients. *Clin Experiment Ophthalmol*. 2004 Feb;32(1):62-6. [Article]
18. Thakur KK, Gupta KK, Mishra SK, Jha N, Chaudhary S, Singh AK. Assessment of Ocular Surface Disease Status in New Patients Initiating Anti-Glaucoma Medication: A Prospective Study. *Birat J Health Sci [Internet]*. 2025 Dec 31 [cited 2026 Feb 12];10(3):60-4. [Article]
19. Thirunavukkarasu A, Karunakaran R. A Study on Evaluation of Ocular Surface Disease in Patients with Glaucoma. 2019;7(1).
20. Vijaya Pai H, Hareesh Reddy LS. Prevalence of ocular surface disease in patients with glaucoma on topical medications. *Asian J Ophthalmol [Internet]*. 2018 Jan 1 [cited 2026 Feb 12];16(2):101-9. [Article]
21. Jaenen N, Baudouin C, Pouliquen P, Manni G, Figueiredo A, Zeyen T. Ocular Symptoms and Signs with Preserved and Preservative-Free Glaucoma Medications. *Eur J Ophthalmol [Internet]*. 2007 May 1 [cited 2026 Jan 10];17(3):341-9. doi: <https://doi.org/10.1177/112067210701700311>
22. Rossi GCM, Pasinetti GM, Scudeller L, Bianchi PE. Ocular surface disease and glaucoma: how to evaluate impact on quality of life. *J Ocul Pharmacol Ther Off J Assoc Ocul Pharmacol Ther*. 2013 May;29(4):390-4. [Article]
23. Zhu W, Xu J. Age-Related Changes of Human Conjunctiva on in vivo Confocal Microscopy. *Invest Ophthalmol Vis Sci [Internet]*. 2009 Apr 28 [cited 2026 Jan 9];50(13):336-336. [Article]
24. Kim H, An Y, Hwang WJ. Gender differences in dry eye disease symptoms associated with psychological health indicators among adults using mobile mental health apps. *PLOS ONE [Internet]*. 2023 Jan 17 [cited 2026 Jan 10];18(1):e0278921. [Article]
25. Sullivan DA, Rocha EM, Aragona P, Clayton JA, Ding J, Golebiowski B, et al. TFOS DEWS II Sex, Gender, and Hormones Report. *Ocul Surf*. 2017 Jul;15(3):284-333. [ScienceDirect]
26. Hong S, Lee CS, Seo KY, Seong GJ, Hong YJ. Effects of topical antiglaucoma application on conjunctival impression cytology specimens. *Am J Ophthalmol*. 2006 Jul;142(1):185-6. [ScienceDirect]
27. Turaçlı E, Budak K, Kaur A, Mizrak B, Ekinci C. The effects of long-term topical glaucoma medication on conjunctival impression cytology. *Int Ophthalmol [Internet]*. 1997 Jan 1 [cited 2026 Jan 9];21(1):27-33. doi: <https://doi.org/10.1023/A:1005892426045>